Why Is It So Hard to Go to Bed With ADHD?

Many adults with ADHD report difficulty winding down, stopping an interesting activity, or getting themselves through the steps required to go to bed. Sleep problems are also common in ADHD.

Bedtime difficulty can involve time awareness, task switching, stimulation, routines, and sleep biology - not just poor discipline.

What Bedtime Difficulty Can Look Like

- losing track of time at night
- saying ‘one more thing’ for an hour
- feeling tired but not ready to stop
- avoiding the steps of getting ready for bed
- getting a burst of energy late at night
- staying up for private or uninterrupted time

Why ADHD May Contribute

- hyperfocus makes stopping difficult
- weak time awareness makes bedtime feel less urgent
- boring routines are hard to initiate
- evening may finally provide quiet or autonomy
- some people with ADHD have delayed sleep timing or other sleep disorders

Sleep Problems Deserve Attention

ADHD and sleep can worsen each other. Poor sleep can increase inattention, irritability, memory problems, and executive functioning difficulties.

- snoring or breathing pauses
- restless legs
- severe daytime sleepiness
- persistent insomnia
- a very delayed sleep schedule

Those symptoms may need medical evaluation rather than being assumed to be ‘just ADHD.’

What Can Help

Make bedtime easier to enter instead of relying on a final burst of willpower.

Reduce Friction

- set a ‘start getting ready’ alarm, not only a bedtime alarm
- do bedtime steps earlier when possible
- make the routine visible and short
- lower lights and stimulation before bed
- use app limits or device charging outside the bedroom if helpful

Support Your Sleep Rhythm

- keep wake time reasonably consistent
- get morning light exposure
- avoid using late-night urgency as your only personal time
- talk with a clinician if sleep problems are persistent or severe

Medication timing, caffeine, mood symptoms, and other health conditions can also affect sleep.

The Main Point

Difficulty going to bed can be an executive functioning and sleep-regulation problem. Better sleep support can improve attention and emotional regulation the next day.

Sources: National Institute of Mental Health, “Attention-Deficit/Hyperactivity Disorder (ADHD)”; NICE, *Attention deficit hyperactivity disorder: diagnosis and management* (NG87). European Consensus Statement on diagnosis and treatment of adult ADHD; reviews of sleep and ADHD.

Prepared by Flourishing Women LLC.